


## Ch 5. Perception and Encoding

Cognitive Neuroscience: The Biology of the Mind, 2<sup>nd</sup> Ed.,  
M. S. Gazzaniga, R. B. Ivry, and G. R. Mangun, Norton, 2002.

Summarized by  
Y.-J. Park, M.-H. Kim, and B.-T. Zhang  
Biointelligence Laboratory, Seoul National University  
<http://bi.snu.ac.kr/>

1

### Disorders of Perception: A Case Study



© 2006, SNU CSE BioIntelligence Lab, <http://bi.snu.ac.kr/>

2

# Contents

- Overview of Neural Pathways
- Parallel Processing in the Visual System
- Cortical Visual Areas
- Deficits in Feature Perception
- Independent or Convergent Pathways
- Dissociations of Cortical and Subcortical Visual Pathways
- Auditory Perception
- Summary

© 2006, SNU CSE BioIntelligence Lab, <http://bi.snu.ac.kr/>

3


## Overview of Neural Pathways

4

## Overview of Neural Pathways

### ● The Eye, Retina, and Receptors

- ◆ As light passes through the lens of eye, the image is inverted and focus on the retina
- ◆ Rods are sensitive to low levels of **light** stimulation at night
- ◆ Cones are most active during daytime vision and essential for **color**
- ◆ Rods and Cones have differential distribution




© 2006, SNU CSE BioIntelligence Lab, <http://bi.snu.ac.kr/>

5

## Overview of Neural Pathways

### ● The Eye, Retina, and Receptors




- ◆ Spectral sensitivity functions for rods and the three types of cone

© 2006, SNU CSE BioIntelligence Lab, <http://bi.snu.ac.kr/>

6

## Overview of Neural Pathways



- From the Eye to the Central Nervous System

- ◆ The optic fibers from the temporal half of the retina project ipsilaterally, while the nasal fibers cross over at the optic chiasm
- ◆ In this way, the input from each visual field is projected to the **primary visual cortex** in the contralateral hemisphere after the fibers synapse in the **lateral geniculate nucleus (LGN)** 90% (geniculo-cortical pathway)
- ◆ A small percentage 10% of visual fibers of the optic nerve terminate in the **superior colliculus and pulvinar nucleus**

© 2006, SNU CSE BioIntelligence Lab, <http://bi.snu.ac.kr/>

7


## Parallel Processing in the Visual System

8

## Parallel Processing in the Visual System

- Organization of the Lateral Geniculate Nucleus (LGN)

- ◆ The LGN contains six well-defined layers
- ◆ Three of the layers receive input from one retina while the other three receive input from the other retina
- ◆ The multilayered system is not simply redundant




© 2006, SNU CSE BioIntelligence Lab, <http://bi.snu.ac.kr/>

9

## Parallel Processing in the Visual System

- Organization of the Lateral Geniculate Nucleus (LGN)

- ◆ The bottom two layers are referred to as the magnocellular system (M system)
- ◆ Smaller cells in the upper four layers constitute the parvocellular system (P System)
- ◆ 80% of the LGN neurons are part of the P system




© 2006, SNU CSE BioIntelligence Lab, <http://bi.snu.ac.kr/>

10

## Parallel Processing in the Visual System

- Multiple Pathways in the Visual Cortex

- ◆ Primary visual cortex (or Striate Cortex)
- ◆ It is located medially and buried below the superficial surface of the cortex along the calcarine sulcus




© 2006, SNU CSE BioIntelligence Lab, <http://bi.snu.ac.kr/>

11

## Parallel Processing in the Visual System

- Multiple Pathways in the Visual Cortex

- ◆ The segregation of M and P pathways is maintained in the cortex
- ◆ Axons from both regions terminate in layer 4 of the striate cortex
- ◆ While all of the inputs terminate in layer 4, the parvocellular inputs synapse on intracortical neurons that terminate in layers 2 and 3




12

# Cortical Visual Areas

13


## Cortical Visual Areas



© 2006, SNU CSE BioIntelligence Lab, <http://bi.snu.ac.kr/>

14

## Cortical Visual Areas



- ◆ Directional and speed tuning of a neuron from area MT
- ◆ Top: A rectangle was moved through the receptive field of this cell in various directions
- ◆ Bottom: The graph shows speed tuning for a cell in MT

© 2006, SNU CSE BioIntelligence Lab, <http://bi.snu.ac.kr/>

15

## Cortical Visual Areas

### ● Cellular Correlates of Visual Features

- ◆ This table summarizes stimulus variations used in neurophysiological studies to identify the representational characteristics of cells in the M, P-blob, and P-interblob pathways
- ◆ Neurons in the M pathway are movement sensitive and color insensitive
- ◆ In contrast, neurons in the P-blob pathway are highly selective to color and are minimally responsive to movement or changes in orientation

Parallel Pathways in Visual Perception			
Neural Structure	Cell Types		
<b>Thalamus (LGN)</b>	<b>Magnocellular</b>	<b>Parvocellular</b>	
Area 17 (V1)	Layer 4b	Blobs	Interblobs
Area 18 (V2)	Thick stripes	Thin stripes	Interstripes
Cellular Correlates			
Contrast (brightness)	high	high	low
Location	low	low	high
Motion	high	low	middle
Color	low	high	middle
Orientation	middle	low	high


© 2006, SNU CSE BioIntelligence Lab, <http://bi.snu.ac.kr/>

16



## Cortical Visual Areas

- Imaging Visual Areas in Humans
  - ◆ Positron Emission Tomography (PET)




© 2006, SNU CSE BioIntelligence Lab, <http://bi.snu.ac.kr/>

17

## Cortical Visual Areas

- Imaging Visual Areas in Humans
  - ◆ The neural basis of visual illusions
  - ◆ (a) Illusory motion is perceived when one is viewing the Enigma pattern, and activation is observed in area MT
  - ◆ (b) Color aftereffects are produced after staring at the bright green patch
  - ◆ (c) Activation in a visual region anterior to V4 is shown graphically




© 2006, SNU CSE BioIntelligence Lab, <http://bi.snu.ac.kr/>

18

## Cortical Visual Areas

- Analysis and Representation of Visual Features

- ◆ Visual Search




© 2006, SNU CSE BioIntelligence Lab, <http://bi.snu.ac.kr/>

19

## Cortical Visual Areas

- Analysis and Representation of Visual Features

- ◆ Illusions and Feature Independence




## Deficits in Feature Perception

21

## Deficits in Feature Perception

### ● Deficits in Color Perception: Achromatopsia

- ◆ Achromatopsia are disorders of color perception that arise from disturbances of the central nervous system
- ◆ MRI scans showing a small lesion encompassing V4
- ◆ Color perception thresholds in each visual quadrant




© 2006, SNU CSE BioIntelligence Lab, <http://bi.snu.ac.kr/>

22

## Deficits in Feature Perception

### ● Deficits in Color Perception: Achromatopsia



- ◆ Stimuli used to assess form perception in the patient with damage to area V4.
- ◆ On basic tests of luminance (a), orientation (b), and motion (c), the patient's perceptual threshold was similar in all four quadrants.
- ◆ Thresholds for illusory contours (d), and complex shapes (e) were elevated in the upper left quadrant.

© 2006, SNU CSE BioIntelligence Lab, <http://bi.snu.ac.kr/>

23

## Independent or Convergent Pathways

24


## Independent or Convergent Pathways

- The visual system contains multiple pathways, each specialized to abstract specific information.
- Yet the outputs from these systems are designed to complement each other.
- Lesion studies of animals have brought into question the notion that the processing of feature like depth, color, and orientation depends solely on a single pathway.
- Prefer the term **concurrent processing**

## Dissociations of Cortical and Subcortical Visual Pathways

## Dissociations of Cortical and Subcortical Visual Pathways

- Spatial Orientation and Object Perception in the Hamster
  - ◆ To turn their head in the direction of a sunflower seed and to run down a maze and to enter the door behind which a sunflower seed was hidden
  - ◆ This study provide compelling evidence for dissociable functions of the hamsters' superior colliculus and visual cortex



© 2006, SNU CSE BioIntelligence Lab, <http://bi.snu.ac.kr/>


27

## Auditory Perception

28

## Auditory Perception

### ● Overview of the Auditory Pathways




© 2006, SNU CSE BioIntelligence Lab, <http://bi.snu.ac.kr/>

29

## Auditory Perception

### ● Overview of the Auditory Pathways

- ◆ Like as photoreceptors, hair cells of the **cochlea** are the primary receptors.
- ◆ Receptive field of hair cells refer to a coding of sound frequency
- ◆ Hair cells at the thick end, or base of the cochlea, are activated by high-frequency sound; cells at the opposite end, or apex, are activated by low-frequency sound




© 2006, SNU CSE BioIntelligence Lab, <http://bi.snu.ac.kr/>

30

## Auditory Perception

### ● Overview of the Auditory Pathways

- ◆ Cells in one region of an auditory area will respond to low-frequency stimuli; cells in another region will respond to middle or high-frequencies
- ◆ Tonotopic representation revealed with fMRI
- ◆ In most cases, the region responding to the low tones (blue) is more posterior and medial to the region responding to the high tones (red)




© 2006, SNU CSE BioIntelligence Lab, <http://bi.snu.ac.kr/>

31

## Auditory Perception

### ● Concurrent Processing for Sound Localization

- ◆ Barn owls rely on two cues to localize sounds: **interaural time**, and **the difference in the sound's intensity at the two ears**
- ◆ These two cues are processed by independent neural pathways: each cochlear nucleus is composed of two parts, the magnocellular nucleus and the angular nucleus



© 2006, SNU CSE BioIntelligence Lab, <http://bi.snu.ac.kr/>

32



## Summary

33

## Summary

- This chapter provided an overview of the organization of the pathways involved in visual and auditory perception.
- A point emphasized in this chapter is that specialized mechanisms for solving different computational problems have evolved in the brain.

## Thought Questions

- You watch a short video segment in which a large purple dinosaur appears briefly in the left visual field. Trace the flow of information about this stimulus and its separate features(color, shape, luminance, motion, position) from the eye through the secondary visual areas
- Compare and contrast the functional organization of the visual and auditory systems. What are the computational problems that each system must solve, and how are these solutions achieved in the nervous system?

## Thought Questions

- A person arrives at the hospital in a confused state and appears to have some impairments in visual perception. As the attending neurologist, you suspect the person may have had a stroke. How would you go about examining the patient to determine at what level in the visual pathways the damage has occurred? You should emphasize the behavioral tests you would administer, although you should feel free to make predictions about what you expect to see on MRI scans

## Thought Questions

- Define the physiological concepts of the receptive field and visual area. How is the receptive field of a cell established? How are the boundaries between visual areas identified? Can either receptive fields or visual areas be studied noninvasively in humans?
- Much of the focus in this chapter has been on salient visual properties such as color, shape, and motion. In looking around the environment, do these properties seem to reflect the most important cues for a highly skilled visual creature? What other sources of information might an adaptive visual system exploit?